Serial No.: 09/868,884 Filed: February 5, 2002

Page : 2 of 17

Amendments to the Claims:

This listing of claims replaces all prior versions and listings of claims in the application:

Listing of Claims:

1. (Previously presented) A compound of formula (I)

A represents thiophene;

 R^1 represents a phenyl group; said phenyl being optionally substituted by one or more substituents selected independently from halogen, cyano, nitro, -NR³R⁴, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -S(O)_mR¹⁰, -S(O)₂NR⁵R⁶, -NR⁸SO₂R¹⁰, C₁-C₆ alkyl, trifluoromethyl, -(CH₂)_nR¹¹, -O(CH₂)_nR¹¹ or -OR¹²;

 R^2 represents hydrogen, halogen, cyano, nitro, $-NR^{13}R^{14}$, $-CONR^{15}R^{16}$, $-COOR^{17}$, $-NR^{18}COR^{19}$, $-S(O)_mR^{20}$, $-S(O)_2NR^{15}R^{16}$, $-NR^{18}SO_2R^{20}$, C_1 - C_2 alkyl, trifluoromethyl, C_2 - C_3 alkenyl, C_2 - C_3 alkynyl, trifluoromethoxy, C_1 - C_2 alkoxy or C_1 - C_2 alkanoyl;

X represents oxygen or sulfur;

each of R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} and R^{12} independently represent a hydrogen atom or C_1 - C_6 alkyl;

Serial No. : 09/868,884 Filed : February 5, 2002

Page : 3 of 17

 R^{11} represents $NR^{21}R^{22}$ where R^{21} and R^{22} are independently hydrogen or C_1 - C_6 alkyl optionally substituted by C_1 - C_4 alkoxy; or R^{21} and R^{22} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR^{23} group where R^{23} is hydrogen or C_1 - C_6 alkyl; or R^{11} represents OR^{24} where R^{24} represents C_1 - C_6 alkyl;

each of R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} independently represent a hydrogen atom or C_1 - C_2 alkyl;

m represents an integer 0, 1 or 2;

n represents an integer 2, 3 or 4;

and optical isomers, racemates, and tautomers thereof and pharmaceutically acceptable salts or solvates thereof:

provided that:

when A represents thiophene, then R¹ is not 4-pyridinyl or 3-pyrazolyl.

- 2. (Original) A compound of formula (I), according to Claim 1, wherein X represents oxygen.
- 3. (Previously presented) A compound of formula (I), according to Claim 1, in which the group A is substituted as shown below in formula (Ia), where B and D are selected from CR^2 and S, where R^2 is as defined in Claim 1 and R^{25} is hydrogen or C_1 - C_6 alkyl:

$$X = \begin{array}{c} NH_2 \\ NH \\ B \\ O \\ NH_2 \end{array}$$
 (Ia)

4. (Cancelled)

Serial No.: 09/868,884 Filed: February 5, 2002

Page : 4 of 17

- 5. (Cancelled)
- 6. (Previously presented) A compound according to claim 1 in which R² represents H or methyl.
- 7. (Original) A compound according to Claim 6 in which R² represents H.
- 8. (Original) A compound of formula (I), according to claim 1, selected from:
- 3-[(aminocarbonyl)amino]-5-phenyl-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-fluorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-isobutylphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(2-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{4-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-2-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-phenyl-3-thiophenecarboxamide;
- $3-[(aminocarbonyl)amino]-5-\{4-[2-(1-morpholinyl)ethoxy]phenyl\}-2-thiophenecarboxamide;\\$
- 3-[(aminocarbonyl)amino]-5-{4-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;

Serial No.: 09/868,884 Filed: February 5, 2002

Page : 5 of 17

3-[(aminocarbonyl)amino]-5-{4-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{4-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{3-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{3-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{3-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{3-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{3-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{2-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{2-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{2-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide; 3-[(aminocarbonyl)amino]-5-{2-[3-(dimethylamino)propoxylphenyl}-2-thiophenecarboxamide: 2-[(aminocarbonyl)amino]-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methylphenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-ethyl-5-phenyl-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methoxyphenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(4-fluorophenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(3-fluorophenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methoxyphenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(3-chloro-4-methoxyphenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(2-chlorophenyl)-3-thiophenecarboxamide; 2-[(aminocarbonyl)amino]-4-methyl-5-(3-trifluoromethylphenyl)-3-thiophenecarboxamide;

Serial No.: 09/868,884 Filed: February 5, 2002

Page : 6 of 17

2-[(aminocarbonyl)amino]-4-methyl-5-(3-methyl-4-methoxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(3,5-dimethoxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(2,3-dimethoxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(4-isopropylphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(3,4,5-trimethoxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(3,4-dichlorophenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(4-cyanophenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-

thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-4-methyl-5-phenyl-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-cyanophenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-trifluoromethylphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(2,4-difluorophenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-methanesulphonylphenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-thiophenecarboxamide;

Serial No.: 09/868,884 Filed: February 5, 2002

Page : 7 of 17

2-[(aminocarbonyl)amino]-5-(4-[2-(1-(2,2,6,6-tetramethyl)piperidinyl)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-(thiazol-4-yl-methoxy)phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(dimethylamino)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-thiophenecarboxamide:

2-[(aminocarbonyl)amino]-5-(4-[2-(1-morpholinyl)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminothiocarbonyl)amino-5-phenyl-3-thiophenecarboxamide;

and pharmaceutically acceptable salts and solvates thereof.

- 9. (Currently amended) A process for the preparation of a <u>first</u> compound of formula (I), according to claim 1, which comprises:
- (a) reaction of a compound of formula (II):

wherein A, R^1 and R^2 are as defined in Claim 1, with an isocyanate (X = O) or an isothiocyanate (X = S), to produce the first compound of formula (I); or

(b) reaction of compound of formula (III) with a compound of formula (IV)

Serial No. : 09/868,884 Filed : February 5, 2002

Page : 8 of 17

wherein A, X, R¹ and R² are as defined in Claim 1, and LG represents a leaving group, to produce the first compound of formula (I); or

(c) reaction of compound of formula (V) with a compound of formula (VI)

$$R^{1}$$
-LG
$$\begin{array}{c} & & & \\ & &$$

wherein A, X, R¹ and R² are as defined in Claim 1, and LG represents a leaving group, to produce the first compound of formula (I);

and where necessary converting the resultant compound of formula (I), or another salt thereof, into a pharmaceutically acceptable salt thereof; or converting the resultant compound of formula (I) into a further compound of formula (I); and where desired converting the resultant compound of formula (I) into an optical isomer thereof.

Serial No. : 09/868,884

Filed: February 5, 2002

Page : 9 of 17

10. (Previously presented) A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1, in

association with a pharmaceutically acceptable adjuvant, diluent or carrier.

11. (Previously presented) A process for the preparation of a pharmaceutical composition

which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or

solvate thereof, as claimed in claim 1 with a pharmaceutically acceptable adjuvant, diluent or

carrier.

12-19. (Cancelled)

20. (Previously presented) A method of treating an IKK2 mediated disease which comprises

administering to a patient a therapeutically effective amount of a compound of formula (I), or a

pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1.

21. (Currently amended) A method of treating an inflammatory disease, or a disease with an

inflammatory component, in a patient suffering from, or at risk of, said disease, which comprises

administering to the patient a therapeutically effective amount of a compound of formula (I), or a

pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1.

22. (Original) A method according to claim 21, wherein the disease is asthma.

23. (Original) A method according to claim 21, wherein the disease is rheumatoid arthritis.

24. (Original) A method according to claim 21, wherein the disease is multiple sclerosis.

25. (Original) A method according to claim 21, wherein the disease is chronic obstructive

pulmonary disease.

Serial No.: 09/868,884 Filed: February 5, 2002

Page : 10 of 17

26. (Previously presented) A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 8, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

- 27. (New) A process of claim 9, further comprising converting the first compound of formula (I), or a salt thereof, into a pharmaceutically acceptable salt thereof; or converting the first compound of formula (I) into a second compound of formula (I).
- 28. (New) A process of claim 9, further comprising converting the first compound of formula (I) into an optical isomer thereof.
- 29. (New) A method of claim 21, wherein the disease is rhinitis.